

## UNIT 7 – RIGHT ANGLE TRIANGLES

Assignment	Title	Work to complete	Complete
1	<i>Triangles</i>	Labelling Triangles	
2	<i>Pythagorean Theorem</i>	Pythagorean Theorem	
3	<i>Pythagorean Triples</i>	Pythagorean Triples	
	<i>Quiz 1</i>		
4	<i>Trigonometry</i>	Trigonometry	
5	<i>The Sine Ratio The Cosine Ratio The Tangent Ratio</i>	The Trigonometric Ratios	
6	<i>Using The Sine Ratio Using The Cosine Ratio Using The Tangent Ratio Angle of Elevation and Depression</i>	Finding Sides In Right Triangles	
	<i>Quiz 2</i>		
7	<i>Finding Angles in Right Triangles</i>	Finding Angles In Right Triangles	
<b>Practice Test</b>	<b>Practice Test</b> How are you doing?	Get this page from your teacher	
<b>Self-Assessment</b>	<b>Self-Assessment</b> <i>“Traffic Lights”</i>	On the next page, complete the self-assessment assignment.	
<b>Chapter Test</b>	<b>Chapter Test</b> Show me your stuff!		
<b>Mental Math</b>	<b>Mental Math</b> Non-calculator practice		

## Traffic Lights

In the following chart, show how confident you feel about each statement by drawing one of the following: 😊, 😐, or ☹️. Then discuss this with your teacher **BEFORE** you write the test!

Statement	😊 😐 ☹️
After completing this chapter;	
<ul style="list-style-type: none"> <li>I can use the Pythagorean theorem to calculate the missing side of a right triangle</li> </ul>	
<ul style="list-style-type: none"> <li>I know when to choose sine (sin), cosine (cos) or tangent(tan) based on the information given</li> </ul>	
<ul style="list-style-type: none"> <li>I can use the three basic trigonometric functions (sin, cos, tan) to find a missing side or angle of a right triangle</li> </ul>	
<ul style="list-style-type: none"> <li>I can determine the angle of elevation and the angle of depression for words or a diagram, and use them with the trigonometric ratios</li> </ul>	
<ul style="list-style-type: none"> <li>I can determine places in the workplace where I could use trigonometry</li> </ul>	

### Vocabulary: Unit 7

angle of depression

angle of elevation

cosine

hypotenuse

leg

Pythagorean Theorem

Pythagorean Triple

right triangle

sine

tangent


## **TRIANGLES**

In this unit, you will be looking at triangles, specifically right angle triangles, also called right triangles. You will learn about Pythagorean Theorem and the basic trigonometric ratios. But first it is necessary to start with some facts about triangles.


Fact 1: Every triangle contains 3 sides and 3 angles or vertices (plural of vertex).

Fact 2: The measurements of these angles always total  $180^{\circ}$ . Remember this from the last unit??

Fact 3: To identify the side or vertex in a triangle, it is important to label the triangle following a standard routine. Each vertex of a triangle is labeled with a capital case letter – like “A” - and each side is labeled with the lower case letter that matches the opposite vertex. An example is below.



Another way to label the sides is with the capital letters of the two vertices the side connects. An example is below.




Side  $a$  can be called BC.  
Side  $b$  can be called AC.  
Side  $c$  can be called AB.


## ASSIGNMENT 1 – LABELLING TRIANGLES

1) Label each side of the triangles below using a single lower case letter matching the opposite vertex.


a)




b)



c)




d)




2) Label each vertex of the triangles below using a single capital letter matching the opposite side.


a)




b)



c)




d)




## **PYTHAGOREAN THEOREM**

Pythagorean Theorem states the relationship between the sides of a right triangle. So, more facts about triangles are necessary.

Fact 4: A triangle that contains a  $90^{\circ}$  angle (a right angle) is called a right triangle (or right-angle triangle).



Fact 5: The side of the triangle that is opposite the  $90^{\circ}$  angle is always called the **hypotenuse**. It is labelled in the triangle below. The other two sides of the triangle are called legs.




Fact 6: The hypotenuse is always the longest side in the triangle. It is always opposite the largest angle which is the  $90^{\circ}$  or right angle.


Fact 7: Pythagorean Theorem states that in any right triangle, the sum of the squares of the lengths of the legs is equal to the square of the length of the hypotenuse. So in  $\Delta ABC$  with the right angle at C, the following relationship is true:

$$c^2 = a^2 + b^2$$

where a and b are the other 2 legs of the triangle.



Often Pythagorean Theorem is illustrated as the square of the sides as follows:



Notice that the length of side  $a$  is 3 boxes, side  $b$  is 4 boxes, and side  $c$  is 5 boxes. So if we calculate the area of each square, the following is true:

$$\begin{aligned} c \times c &= c^2 = 5 \times 5 = 25 \\ b \times b &= b^2 = 4 \times 4 = 16 \\ a \times a &= a^2 = 3 \times 3 = 9 \end{aligned}$$

And we know that

$$c^2 = a^2 + b^2$$

So  $25 = 9 + 16$  which is a true statement!

We can also rearrange the equation to find the length one of the legs;


$$c^2 = a^2 + b^2$$

$$a^2 = c^2 - b^2$$

$$b^2 = c^2 - a^2$$

When we use Pythagorean Theorem to find a length of the hypotenuse or a leg, you need to have a calculator that has the square root function  $\sqrt{\quad}$  on it. The computer symbol looks like this:  $\sqrt{\quad}$  or  $\sqrt{\quad}$

Example 1: Use Pythagorean Theorem to find the length of the missing side to one decimal place.




Solution:

$$\begin{aligned} q^2 &= p^2 + r^2 \\ q^2 &= 5.2^2 + 3.8^2 \\ q^2 &= 27.04 + 14.44 \\ q^2 &= 41.48 \\ \sqrt{q^2} &= \sqrt{41.48} \\ q &\approx 6.44 \text{ cm} \end{aligned}$$

Side  $q$  is approximately 6.4 cm

Example 2: Use Pythagorean Theorem to find the length of the missing side to one decimal place.




Solution:  $c^2 = a^2 + b^2$   
 So,  $b^2 = c^2 - a^2$   
 $b^2 = 12.8^2 - 10.78^2$   
 $b^2 = 163.84 - 116.21$   
 $b^2 = 47.66$   
 $\sqrt{b^2} = \sqrt{47.66}$   
 $b \approx 6.90$  cm  
 Side b is approximately 6.9 cm


**ASSIGNMENT 2 – PYTHAGOREAN THEOREM**

1) Using the following triangles, use lettering provided to state the Pythagorean relations that apply.

a)



b)



2) Find the missing value in each of the following to 2 decimal places.

a)  $p^2 = 6^2 + 9^2$


b)  $m^2 = 4^2 + 7^2$

c)  $y^2 = 8^2 - 5^2$


d)  $z^2 = 10^2 - 5^2$

3) Calculate the missing side length to 1 decimal place as needed.


a)




b)




c)




d)




4) Find the length of the bracket in the picture, to one decimal place.




5) What is the length of the guy wire in the picture below, to one decimal place?



6) A ramp into a house rises up 3.5 meters over a horizontal distance of 10.5 meters. How long is the ramp? Draw a diagram and show your work.



- 7) You need to find the width of a lake,  $PQ$ , as shown. The measurements of the other sides are given on the diagram. You are certain that  $\angle P = 90^\circ$ . What is the width of the lake?




- 8) A ladder is leaned against a house. The base of the ladder is  $d$  feet away from the house. Draw a diagram and then write the Pythagorean relationship that exists for these lengths. Use  $l$  for the ladder,  $h$  for the house, and  $d$  for the distance the ladder is from the house. You are not required to solve this question.

- 9) A 40 foot ladder reaches 38 feet up the side of a house. How far from the side of the house is the base of the ladder? Draw a diagram and show your work.

- 10) A flagpole is 12 metres tall. It makes a shadow on the ground that is 15 metres long. How long is a line that joins the top of the flagpole with the end of the shadow? Draw a diagram and show your work.

## PYTHAGOREAN TRIPLES

A **Pythagorean Triple** is a set of three numbers that satisfy the Pythagorean Theorem and are all whole numbers (no decimals). If a set of numbers satisfies the Pythagorean relationship, then the triangle must be a right triangle. An example is shown below.



$$\begin{aligned}\text{Left Side} &= c^2 \\ &= 13^2 \\ &= \mathbf{169}\end{aligned}$$


$$\begin{aligned}\text{Right Side} &= a^2 + b^2 \\ &= 12^2 + 5^2 \\ &= 144 + 25 \\ &= \mathbf{169}\end{aligned}$$

Therefore, Left Side = Right Side. This triangle is a right angle triangle, and the set of numbers, 5, 12, and 13 are a Pythagorean Triple.


## ASSIGNMENT 3 – PYTHAGOREAN TRIPLES

1) Which of the following triangles are right triangles? Show your work as proof.


a)




b)



c)



d)



**ASK YOUR TEACHER FOR UNIT QUIZ 1.**


## **TRIGONOMETRY**

Trigonometry is one of the most important topics in mathematics. Trigonometry is used in many fields including engineering, architecture, surveying, aviation, navigation, carpentry, forestry, and computer graphics. Also, until satellites, the most accurate maps were constructed using trigonometry.

The word **trigonometry** means triangle measurements. It is necessary to finish our triangle facts here.


**Fact 8:** In trigonometry, the other two sides (or legs) of the triangle are referred to as the **opposite** and **adjacent** sides, depending on their relationship to the angle of interest in the triangle.

In this example, if we pick angle DEF – the angle labelled with the Greek letter  $\theta$  – then we are able to distinguish the sides as illustrated in the diagram below.



The side that is opposite the angle of interest, in this case  $\theta$ , is called the **opposite** side. The side that is nearest to angle  $\theta$  and makes up part of the angle is called the **adjacent** side. To help you, remember that adjacent means beside. Although the hypotenuse occupies one of the two adjacent positions, it is **never** called the adjacent side. It simply remains the hypotenuse. This is why it is identified first. It is recommended to label the side in the order hypotenuse, opposite, and finally adjacent. You may use initials for these side, h, o, and a, but always use lower case letters to avoid mixing up the labelling with a vertex.

Example 1: Using the triangle below, answer the questions.




- 1) What is the hypotenuse? \_\_\_\_\_
- 2) What is the opposite side to  $\theta$ ? \_\_\_\_\_
- 3) What is the adjacent side to  $\theta$ ? \_\_\_\_\_

Solution:

- 1) What is the hypotenuse? 15
- 2) What is the opposite side to  $\theta$ ? 9
- 3) What is the adjacent side to  $\theta$ ? 12

This example uses the same triangle as in Example 1; however, this time, the other acute angle is labelled as  $\theta$ . This is done to show that the opposite and adjacent sides switch when the other angle is the angle of interest. The hypotenuse **always** stays the same.

Example 2: Using the triangle below, answer the questions.



- 1) What is the hypotenuse? \_\_\_\_\_
- 2) What is the opposite side to  $\theta$ ? \_\_\_\_\_
- 3) What is the adjacent side to  $\theta$ ? \_\_\_\_\_


Solution:

- 1) What is the hypotenuse? 15
- 2) What is the opposite side to  $\theta$ ? 12
- 3) What is the adjacent side to  $\theta$ ? 9

## ASSIGNMENT 4 – TRIGONOMETRY


For each of the right triangles below, mark the hypotenuse, and the sides that are opposite and adjacent sides to  $\theta$  as shown in the example.

Example:




h = hypotenuse  
o = opposite  
a = adjacent


1)




2)



3)



4)



## TRIGONOMETRIC RATIOS

In the previous unit about similar figures, you learned that the ratios of corresponding sides of similar triangles are equal. When the angles of different triangles are the same, the ratio of the sides within the triangle will always be the same. They depend only on the measure of the angle of interest, not the size of the triangle. These ratios are the trigonometric ratios.


There are three trigonometric ratios we are concerned with: sine, cosine, and tangent.

### THE SINE RATIO

The *sine of angle  $\theta$*  means the ratio of the length of opposite side to the length of the hypotenuse. It is abbreviated as **sin  $\theta$**  but read as sine  $\theta$ . It is written like this:

$$\sin \theta = \frac{\textit{opposite}}{\textit{hypotenuse}} \quad \text{or} \quad \sin \theta = \frac{o}{h}$$

Example 1: Find the sine of  $\theta$  in this triangle. Round to 4 decimal places.



Solution:

The opposite side is 5 and the hypotenuse is 13. So

$$\sin \theta = \frac{o}{h} = \frac{5}{13} = 0.3846 \quad \text{So} \quad \sin \theta = 0.3846$$

Note: Rounding to 4 decimal places is standard when calculating trigonometric ratios.

Example 2: Use your calculator to determine the following sine ratios. Round to 4 decimal places.

- a)  $\sin 15^\circ$                       b)  $\sin 67^\circ$                       c)  $\sin 42^\circ$

**\*\*\*\*\* REMEMBER TO SET YOUR CALCULATOR ON DEGREES (DEG) \*\*\*\*\***

Solution: Type “sin” followed by the angle, and then “=” to solve


- a)  $\sin 15^\circ = 0.2588$       b)  $\sin 67^\circ = 0.9205$                       c)  $\sin 42^\circ = 0.6691$

## THE COSINE RATIO

The *cosine of angle  $\theta$*  means the ratio of the adjacent side to the hypotenuse. It is abbreviated as **cos  $\theta$**  but read as cosine  $\theta$ . It is written like this:

$$\cos \theta = \frac{\textit{adjacent}}{\textit{hypotenuse}} \quad \text{or} \quad \cos \theta = \frac{a}{h}$$

Example 1: Find the cosine of  $\theta$  in this triangle.



Solution:

The adjacent side is 12 and the hypotenuse is 13. So

$$\cos \theta = \frac{a}{h} = \frac{12}{13} = 0.9231$$

Note: Rounding to 4 decimal places is standard when calculating trigonometric ratios.

Example 2: Use your calculator to determine the following cosine ratios. Round to 4 decimal places.

- a)  $\cos 15^\circ$                       b)  $\cos 67^\circ$                       c)  $\cos 42^\circ$

**\*\*\*\*\* REMEMBER TO SET YOUR CALCULATOR ON DEGREES (DEG) \*\*\*\*\***

Solution: Type “cos” followed by the angle, and then “=” to solve


- a)  $\cos 15^\circ = 0.9659$     b)  $\cos 67^\circ = 0.3907$                       c)  $\cos 42^\circ = 0.7431$

## THE TANGENT RATIO

The *tangent of angle  $\theta$*  means the ratio of the opposite side to the adjacent side. It is abbreviated as **tan  $\theta$**  but read as tangent  $\theta$ . It is written like this:

$$\tan \theta = \frac{\textit{opposite}}{\textit{adjacent}} \quad \text{or} \quad \tan \theta = \frac{o}{a}$$

Example 1: Find the tangent of  $\theta$  in this triangle.



Solution:

The opposite side is 5 and the adjacent side is 12. So

$$\tan \theta = \frac{o}{a} = \frac{5}{12} = 0.4167$$

Note: Rounding to 4 decimal places is standard when calculating trigonometric ratios.

Example 2: Use your calculator to determine the following tangent ratios. Round to 4 decimal places.

- a)  $\tan 15^\circ$                       b)  $\tan 67^\circ$                       c)  $\tan 42^\circ$

**\*\*\*\*\* REMEMBER TO SET YOUR CALCULATOR ON DEGREES (DEG) \*\*\*\*\***



Solution: Type “tan” followed by the angle, and then “=” to solve

- a)  $\tan 15^\circ = 0.2679$       b)  $\tan 67^\circ = 2.3559$                       c)  $\tan 42^\circ = 0.9004$



## ASSIGNMENT 5 – THE TRIGONOMETRIC RATIOS

1)



a) Calculate the value of **sin X** to two decimal places.



b) Calculate the value of **cos X** to two decimal places.



c) Calculate the value of **tan X** to two decimal places.



2) Use your calculator to determine the value of each of the following sine ratios to four decimal places.

a)  $\sin 10^\circ =$  \_\_\_\_\_

b)  $\sin 48^\circ =$  \_\_\_\_\_

c)  $\sin 77^\circ =$  \_\_\_\_\_

d)  $\sin 85^\circ =$  \_\_\_\_\_

e)  $\cos 10^\circ =$  \_\_\_\_\_

f)  $\cos 48^\circ =$  \_\_\_\_\_

g)  $\cos 77^\circ =$  \_\_\_\_\_

h)  $\cos 85^\circ =$  \_\_\_\_\_

i)  $\tan 10^\circ =$  \_\_\_\_\_

j)  $\tan 48^\circ =$  \_\_\_\_\_

k)  $\tan 77^\circ =$  \_\_\_\_\_

l)  $\tan 85^\circ =$  \_\_\_\_\_

3) There are two special sine ratios. Calculate the following and suggest why the values are what the results give you.

a)  $\sin 0^\circ =$  \_\_\_\_\_

b)  $\sin 90^\circ =$  \_\_\_\_\_

4) There are two special cosine ratios. Calculate the following and suggest why the values are what the results give you.

a)  $\cos 0^\circ =$  \_\_\_\_\_

b)  $\cos 90^\circ =$  \_\_\_\_\_

5) There are some special tangent ratios. Calculate the following and suggest why the values are what the results give you.

a)  $\tan 0^\circ =$  \_\_\_\_\_

b)  $\tan 45^\circ =$  \_\_\_\_\_


c)  $\tan 89^\circ =$  \_\_\_\_\_

d)  $\tan 90^\circ =$  \_\_\_\_\_

## USING THE SINE RATIO


Whenever one side and one angle of a right triangle are already known, the remaining sides can be found using the trigonometric ratios. The sine ratio can be used to find missing parts of a right triangle.

Example 1: Use the sine ratio to find the  $x$  in the triangle below.



Solution:

Step 1: Label the sides of the triangle with **h**, **o** and **a**



Step 2: Circle the number with the side it represents and the unknown ( $x$ ) with the side it represents.

Step 3: Identify the ratio required to solve for  $x$

Since **o** and **h** are being used, the correct ratio is **sin  $\theta$**

Step 4: Substitute the correct values into the correct ratio.

$$\sin \theta = \frac{o}{h}$$

$$\sin 35^\circ = \frac{9}{x}$$

Step 5: Solve using the process Cross Multiply and Divide.



$$\text{Since } \sin 35^\circ = \frac{\sin 35}{1}, \text{ then } \sin 35^\circ = \frac{9}{x} \text{ becomes } \frac{\sin 35}{1} = \frac{9}{x}$$

$$\begin{aligned} x &= 9 \times 1 \div \sin 35^\circ \\ &= 15.7 \text{ m} \end{aligned}$$

Example 2: A ladder 8.5 m long makes an angle of  $72^\circ$  with the ground. How far up the side of a building will the ladder reach?

Solution:

Sketch a diagram and place the information from the question on this diagram. Remember that there will always be a right triangle in your diagram. It is often helpful to draw that triangle and copy the key information from the sketch.



Step 1: Label the sides of the triangle with **h**, **o** and **a**  
See above right.

Step 2: Circle the number with the side it represents and the unknown ( $x$ ) with the side it represents.

Step 3: Identify the ratio required to solve for  $x$

Since **o** and **h** are being used, the correct ratio is **sin  $\theta$**

Step 4: Substitute the correct values into the correct ratio.

$$\sin \theta = \frac{o}{h}$$

$$\sin 72^\circ = \frac{x}{8.5}$$

Step 5: Solve using the process Cross Multiply and Divide.


$$\text{Since } \sin 72^\circ = \frac{\sin 72}{1}, \text{ then } \sin 72^\circ = \frac{x}{8.5} \text{ becomes } \frac{\sin 72}{1} = \frac{x}{8.5}$$

$$\begin{aligned} x &= \sin 72^\circ \times 8.5 \div 1 \\ &= 8.1 \text{ m} \end{aligned}$$

## USING THE COSINE RATIO


Whenever one side and one angle of a right triangle are already known, the remaining sides can be found using the trigonometric ratios. The cosine ratio can be used to find missing parts of a right triangle.

Example 1: Use the correct trig ratio to find the  $x$  in the triangle below.



Solution:

Step 1: Label the sides of the triangle with **h**, **o** and **a**



Step 2: Circle the number with the side it represents and the unknown ( $x$ ) with the side it represents.

Step 3: Identify the ratio required to solve for  $x$

Since **a** and **h** are being used, the correct ratio is **cos  $\theta$**

Step 4: Write down the chosen ratio and substitute the correct values into the correct ratio.

$$\cos \theta = \frac{a}{h}$$

$$\cos 30^\circ = \frac{x}{5}$$

Step 5: Solve using the process Cross Multiply and Divide.


$$\text{Since } \cos 30^\circ = \frac{\cos 30}{1}, \text{ then } \cos 30^\circ = \frac{x}{5} \text{ becomes } \frac{\cos 30}{1} = \frac{x}{5}$$

$$\begin{aligned} x &= \cos 30^\circ \times 5 \div 1 \\ &= 4.3 \text{ cm} \end{aligned}$$

## USING THE TANGENT RATIO


Whenever one side and one angle of a right triangle are already known, the remaining sides can be found using the trigonometric ratios. The tangent ratio can be used to find missing parts of a right triangle.

Example 1: Use the correct trig ratio to find the  $x$  in the triangle below.



Solution:

Step 1: Label the sides of the triangle with **h**, **o** and **a**



Step 2: Circle the number with the side it represents and the unknown ( $x$ ) with the side it represents.

Step 3: Identify the ratio required to solve for  $x$

Since **o** and **a** are being used, the correct ratio is **tan  $\theta$**

Step 4: Substitute the correct values into the correct ratio.

$$\tan \theta = \frac{o}{a}$$

$$\tan 15 = \frac{2}{x}$$


Step 5: Solve using the process Cross Multiply and Divide.

$$\text{Since } \tan 15^\circ = \frac{\tan 15}{1}, \text{ then } \tan 15^\circ = \frac{2}{x} \text{ becomes } \frac{\tan 15}{1} = \frac{2}{x}$$

$$\begin{aligned} x &= 2 \times 1 \div \tan 15^\circ \\ &= 7.5 \text{ mm} \end{aligned}$$

## ANGLE OF ELEVATION AND DEPRESSION

When you look up at an airplane flying overhead for example, the angle between the horizontal and your line of sight is called the angle of elevation.




When you look down from a cliff to a boat passing by, the angle between the horizontal and your line of sight is called the angle of depression.

When you are given the angle of depression, it is important to carefully use this angle in your triangle.

Example 1: You are standing at the top of a cliff. You spot a boat 200 m away at an angle of depression of  $55^\circ$  to the horizon. How far is the boat from the coast? Draw a diagram to illustrate this situation.

Solution: Draw a diagram, label it with the information, and then solve the triangle.




The angle inside the triangle is the complement to the angle of depression. To find that angle, do the following:  
 $\theta = 90^\circ - 55^\circ$   
 $\theta = 35^\circ$


## **ASSIGNMENT 6 – FINDING SIDES IN RIGHT TRIANGLES**

1) Calculate the length of the side indicated in the following diagrams. Round to **one** decimal place.


a)




b)




c)




d)



e)



f)



2) The angle of elevation from the ground to the top of a flagpole is  $14^\circ$ . Draw a sketch to illustrate this situation.


3) From a building, the angle of depression to a fountain is  $62^\circ$ . The fountain is 75 metres away (along the ground) from the building. Draw a sketch to illustrate this situation.

4) From the top of a 45 metre tall oil rig's mast, the angle of depression to the ground is  $12^\circ$ . Draw a sketch to illustrate this situation, and then find the distance from the top of the mast to the ground along the sight line.

5) A ramp makes an angle of  $22^\circ$  with the ground. If the end of the ramp is 1.5 m above the ground, how long is the ramp?

6) A weather balloon, which is blowing in the wind, is tied to the ground with a 15 m string. How high is the balloon if the angle between the string and the ground is  $38^\circ$ ?

7) A child's slide rises to a platform at the top is  $20^\circ$ . If the horizontal distance that the slide covers is 25 m long, how long is the slide?



8) A flagpole is anchored to the ground by a guy wire that is 12 m long. The guy wire makes an angle of  $63^\circ$  with the ground. How far from the base of the flagpole must the guy wire be anchored into the ground?

9) A man stands 15 m from the base of a tree. He views the top of the tree at an angle of elevation of  $58^\circ$ . How tall is the tree?

10) How far from the side of a house is the base of a ladder if the angle of elevation is  $70^\circ$  and the ladder reaches 15 feet up the side of the house?

**ASK YOUR TEACHER FOR UNIT QUIZ 2.**

## **FINDING ANGLES IN RIGHT TRIANGLES**

So far in this unit, you have used the trigonometric ratios to find the length of a side. But if you know the trigonometric ratio, you can calculate the size of the angle. This requires an “inverse” operation. You can use your calculator to find the opposite of the usual ratio provided you can calculate the ratio. To do this you need 2 sides in the triangle. You can think of the inverse in terms of something simpler: addition is the opposite or inverse of subtraction. In the same way, trig functions have an inverse.

To calculate the inverse, you usually use a 2nd function and the sin/cos/tan buttons on your calculator in sequence. If you look at your calculator just above the sin/cos.tan buttons, you should see the following:  $\sin^{-1}$ ,  $\cos^{-1}$ ,  $\tan^{-1}$ . These are the inverse functions. If you use these buttons, you will be able to turn a ratio into an angle.

Example 1: Calculate each angle to the nearest whole degree.

a)  $\sin X = 0.2546$

b)  $\cos Y = 0.1598$

c)  $\tan Z = 3.2785$

Solution: Use the appropriate inverse function on your calculator.


NOTE: Every calculator is different in how the buttons are keyed in order to achieve the desired outcome. Most calculators will need to key “**2ndF sin**” or “**Shift sin**” in order to get  $\sin^{-1}$  displayed. Then key in the value with or without brackets as necessary.

a)  $\sin X = 0.2546$   
 $X = \sin^{-1}(0.2546)$   
 $X = 14.74988^{\circ}$                       Angle X is  $15^{\circ}$ .

b)  $\cos Y = 0.1598$   
 $Y = \cos^{-1}(0.1598)$   
 $Y = 80.8047^{\circ}$                       Angle Y is  $81^{\circ}$ .

c)  $\tan Z = 3.2785$   
 $Z = \tan^{-1}(3.2785)$   
 $Z = 73.03737^{\circ}$                       Angle Z is  $73^{\circ}$ .

Example 2: Determine the angle  $\theta$  in the following triangle.




Solution:

- 1) h, o, a the triangle
- 2) Circle the letters with their partner numbers
- 3) Choose the appropriate trig ratio. In this case, it is tangent.
- 4) Write down the ratio and fill it in.

$$\tan \theta = \frac{o}{a}$$

$$\tan \theta = \frac{5}{3}$$



- 5) Divide the numerator (top number) by the denominator (bottom number) in the fraction to get a decimal number.

$$\tan \theta = 1.66666$$

- 6) Use the inverse function to solve for  $\theta$ .

$$\theta = \tan^{-1}(1.66666)$$

$$\theta = 59.0352^{\circ}$$

Angle  $\theta$  is approximately  $59^{\circ}$ .

## **ASSIGNMENT 7 – FINDING ANGLES IN RIGHT TRIANGLES**

1) Calculate the following angles to the nearest whole degree.

a)  $\sin D = 0.5491$

b)  $\cos F = 0.8964$


c)  $\tan G = 2.3548$

d)  $\sin P = 0.9998$


e)  $\cos Q = 0.3097$

f)  $\tan R = 0.4663$

2) After an hour of flying, a jet has travelled 300 miles, but gone off course 48 miles west of its planned flight path. What angle,  $\theta$ , is the jet off course?




3) At what angle to the ground is an 8 m long conveyor belt if it is fastened 5 m from the base of the loading ramp?



4) If a boat is 150 m from the base of a 90 m cliff, what is the angle of elevation from the boat to the top of the cliff?

5) ) In a right triangle,  $\Delta XYZ$ , the ratio of the opposite side to  $\angle X$  to the hypotenuse is 7:8 or  $\frac{7}{8}$ . What is the approximate size of  $\angle X$ ?

6) What is the angle of depression,  $\theta$ , from the top of a 65 m cliff to an object 48 m from its base?



## **UNIT REVIEW**

The techniques used to solve problems involving right angle triangles have widespread applications in many real-life areas. Often, word problems are presented where Pythagorean Theorem or trigonometry ratios are used to answer a question. The following section of questions requires that you solve these problems in that way. Remember these points to help you solve trig problems:

1. After reading the given information carefully, draw and label a diagram of a triangle if one is not provided.
2. When a side is the unknown and no angle is given, use the Pythagorean Theorem to find the unknown side.
3. Label the triangle with o, a, and h in order to choose the correct trig ratio.
4. When a side is the unknown and an angle is given, use the appropriate trig ratio and only the sin, cos, or tan key on the calculator.
5. When an angle is the unknown, use the 2ndF key in addition to the sin, cos or tan key to get the inverse functions:  $\sin^{-1}$ ,  $\cos^{-1}$ ,  $\tan^{-1}$ .
6. Solve for the unknown.
7. Check to see if your answer is reasonable.

Some students find it helpful to remember the trigonometric relationships of sine, cosine, and tangent with the phrase **SOH CAH TOA** (pronounced sow-a caw toe-a). This comes from the initials from the trig ratios and their sides:

$$\sin = \frac{o}{h} \quad \cos = \frac{a}{h} \quad \tan = \frac{o}{a}$$

You will be given the trigonometric ratios and Pythagorean Theorem in the Data Pages for both your tests and the Provincial exam. You are responsible for knowing how to use and apply these formulas.